

## **CR-SA HYSTERESIS PHENOMENON ON THE BASIS OF NM DATA,**

### **4. COMPARISON OF TIME-LAGS FOR ODD AND EVEN SOLAR CYCLES, RELATIVE ROLE OF DRIFTS.**

L.I. Dorman (1,2), I.V. Dorman (3), N. IUCCI (4), M. Parisi (4), G. VILLORESI (4)

(1) Israel Cosmic Ray Center and Emilio Segre' Observatory, affiliated to Tel Aviv

University, Technion and Israel Space Agency, Israel; (2) IZMIRAN, Russian

Academy of Science, Troitsk, Russia; (3) Institute of History of Science and

Technology, Russian Academy of Science, Moscow, Russia;

(4) Dipartimento di Fisica "E. Amaldi", Università "Roma Tre", Rome, Italy;

Estimations of cosmic ray time-lags in long-term variations for solar cycles 19 and 21 (Dorman et al., 2001a), as well as for cycles 20 and 22 (Dorman et al., 2001b), show that there are large differences between odd and even solar cycles. We assume that this odd-even effect in the hysteresis phenomenon is caused mainly by cosmic ray drift effects. The odd-even effect in time-lags we use for rough estimates of the relative role of drifts in establishing long-term modulation. We found that with changing of assuming relative role of drift effects (amplitude  $A_{dr}$  in %), the time lag ( $X_{max}$  in average months) also changed, but in different directions for odd and even solar cycles: for odd cycles 19, 21  $X_{max}$  decreases with increasing of  $A_{dr}$ , and for even cycles 20, 22  $X_{max}$  increases with increasing of  $A_{dr}$ . We try to use odd-even effect in time-lags for a rough estimation of the relative role of CR drift effects in formation of long-term modulation as well as for estimation of real time lag and dimension of the modulation region. To determine  $X_{max}$  and  $A_{dr}$  separately, we assume that for a first approximation  $X_{max}$  and  $A_{dr}$  are about the same in odd and even solar cycles. In this case the crossing of dependences of  $X_{max}$  from  $A_{dr}$  for odd and even cycles determines the expected values of  $X_{max}$  and  $A_{dr}$ . On the basis of estimated value of  $X_{max}$  we determine the dimension of modulation region (with taking into account the influence of nonlinear processes on the solar wind speed in the outer Heliosphere according to Le Roux and Fichtner, 1997), the radial diffusion coefficient and transport path as well as expected CR intensity out of the modulation region and absolute CR modulation (relative to the CR intensity level in the interstellar space). We determine also the relative role of drifts and convection-diffusion in formation of observed CR modulation in the Heliosphere in dependence of particle rigidity.

#### REFERENCES:

Dorman, L.I., et al, Proc. 27 ICRC, Paper SH... (2001a).

Dorman, L.I., et al., Proc. 27 ICRC, Paper SH... (2001b).

Le Roux, J. A. and H. Fichtner, Ap. J, 477, L115-L118, 1997.