

## ENERGY SPECTRUM ABOVE $3 \times 10^{18}$ EV OBSERVED WITH AGASA

M. Chikawa (2), M. Fukushima (1), N. Hayashida (1), K. Honda (3), N. Inoue (4), K. Kadota (5), F. Kakimoto (5), S. Kakizawa (6), K. Kamata (7), S. Kawaguchi (8), S. Kawakami (9), Y. Kawasaki (10), N. Kawasumi (11), E. Ku-sano (12), A. M. Mahrous (4), K. Mase (1), T. Minagawa (1), Y. Morizane (2), M. Nagano (13), H. Ohoka (1), S. Osone (1), **N. Sakaki** (10), M. Sasaki (1), M. Sasano (14), K. Shinozaki (4), M. Takeda (1), M. Teshima (1), R. Torii (1), I. Tsushima (11), Y. Uchihori (15), T. Yamamoto (1), K. Yasui (2), S. Yoshida (1), H. Yoshii (16) and T. Yoshikoshi (9)

(1) Institute for Cosmic Ray Research, University of Tokyo, Chiba 277-8582, Japan, (2) Department of Physics, Kinki University, Osaka 577-8502, Japan, (3) Faculty of Engineering, Yamanashi University, Kofu 400-8511, Japan, (4) Department of Physics, Saitama University, Urawa 338-8570, Japan, (5) Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan, (6) Faculty of Science, Shinshu University, Matsumoto 390-8621, Japan, (7) Nishina Memorial Foundation, Komagome, Tokyo 113-0021, Japan, (8) Faculty of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan, (9) Department of Physics, Osaka City University, Osaka 558-8585, Japan, (10) RIKEN (The Institute of Physical and Chemical Research), Saitama 351-0198, Japan, (11) Faculty of Education, Yamanashi University, Kofu 400-8510, Japan, (12) KEK, High Energy Accelerator Research Organization, Institute of Particle and Nuclear Studies, Tsukuba 305-0801, Japan, (13) Department of Applied Physics and Chemistry, Fukui University of Technology, Fukui 910-8505, Japan, (14) Communications Research Laboratory, Ministry of Posts and Telecommunications, Tokyo 184-8795, Japan, (15) National Institute of Radiological Sciences, Chiba 263-8555, Japan, (16) Department of Physics, Ehime University, Matsuyama 790-8577, Japan.

[sakaki@icrr.u-tokyo.ac.jp](mailto:sakaki@icrr.u-tokyo.ac.jp)

We report the energy spectrum at energies greater than  $3 \times 10^{18}$  eV using the updated data set of Akeno Giant Air Shower Array (AGASA). In order to increase statistics and cover larger area of the sky, events up to 60 degree in zenith angle are used. This makes the exposure increase by 50% above  $10^{19}$  eV. The energy spectrum of inclined showers agrees with that of vertical showers. The possible origins of highest energy cosmic rays will be discussed.