

UPPER LIMITS OF DIFFUSE GAMMA RAYS FROM THE GALACTIC PLANE USING THE DATA WITH THE TIBET II AND HD ARRAYS

M. Amenomori (1), S. Ayabe (2), S.W. Cui (3), L.K. Ding (3), X.Y. Ding (4), C.F. Feng (5), Z.Y. Feng (6), Y. Fu (5), X.Y. Gao (7), Q.X. Geng (7), H.W. Guo (4), M. He (5), K. Hibino (8), N. Hotta (9), J. Huang (9), Q. Huang (6), A.X. Huo (3), K. Izu (10), H.Y. Jia (6), F. Kajino (11), K. Kasahara (12), Y. Katayose (13), K. Kawata (11), Labaciren (4), G.M. Le (14), J.Y. Li (5), H. Lu (3), S.L. Lu (3), G.X. Luo (3), X.R. Meng (4), K. Mizutani (2), J. Mu (7), H. Nanjo (1), M. Nishizawa (15), **M. Ohnishi** (10), I. Ohta (9), H. Ooura (11), T. Ouchi (8), S. Ozawa (9), J.R. Ren (3), T. Saito (16), M. Sakata (11), T. Sasaki (17), M. Shibata (13), A. Shiomi (10), T. Shirai (8), H. Sugimoto (18), K. Taira (18), M. Takita (10), Y.H. Tan (3), N. Tateyama (8), S. Torii (8), S. Udo (2), T. Utsugi (2), C.R. Wang (5), H. Wang (3), X. Wang (5), X.W. Xu (3,10), L. Xu (5), X.C. Yang (7), Y. Yamamoto (11), Z.H. Ye (14), G.C. Yu (6), A.F. Yuan (4), T. Yuda (19), H.M. Zhang (3), J.L. Zhang (3), N.J. Zhang (5), X.Y. Zhang (5), Zhaxiciren (4) and Zhaxisangzhu (4)

(1) Dept. of Phys., Hirosaki Univ., Hirosaki, Japan, (2) Dept. of Phys., Saitama Univ., Urawa, Japan, (3) Cosmic Ray and HE Astrophys. Lab, IHEP, CAS, Beijing, China, (4) Dept. of Math. and Phys., Tibet Univ., Lhasa, China, (5) Dept. of Phys., Shangdong Univ., Jinan, China, (6) Inst. of Modern Phys., South West Jiaotong Univ., Chengdu, China, (7) Dept. of Phy., Yunnan Univ., Kunming, China, (8) Faculty of Eng., Kanagawa Univ., Yokohama, Japan, (9) Faculty of Ed., Utsunomiya Univ., Utsunomiya, Japan, (10) ICRC, Univ. of Tokyo, Kashiwa, Japan, (11) Dept. of Phys., Konan Univ., Kobe, Japan, (12) Faculty of Systems Eng., Shibaura Inst. of Technology, Omiya, Japan, (13) Dept. of Phys., Yokohama Natl. Univ., Yokohama, Japan, (14) Center of Space Sci. and Application Res., CAS, Beijing, China, (15) Natl. Inst. for Informatics, Tokyo, Japan, (16) Tokyo Metropolitan Coll. of Aeronautical Eng., Tokyo, Japan, (17) CCDL Lab, Waseda Univ., Tokyo, Japan, (18) Shonan Inst. of Technology, Fujisawa, Japan, (19) STEL, Nagoya Univ., Nagoya Japan.
yamamoto@hep.konan-u.ac.jp

The intensity upper limits of 10 TeV region gamma rays from the inner and outer Galactic planes were given using data with the Tibet II and high density (HD) arrays. The present result seems to reject the inverse Compton gamma rays by energetic electrons with a hard power spectrum of -2.0. The data with the HD array at 3 TeV region gives the first upper limits. But, these are not enough to reject both models of the inverse Compton with softer energy spectra and of proton induced gamma rays.