

Momentum spectrum of cosmic muons at a depth of 320 mwe

N.O. Hashim^a, C. Grupen^a, S. Luitz^b, A. Mailov^a, F. Maciuc^c, A.-S. Müller^d,
A. Putzer^e, H.-G. Sander^f, S. Schmeling^g, M. Schmelling^c, R. Tcaciuc^a,
H. Wachsmuth^g, Th. Ziegler^h, K. Zuberⁱ

(a) *Fachbereich Physik, Universität Siegen, D-57068 Siegen, Germany*

(b) *SLAC M/S 41, 2575 Sand Hill Rd, Menlo Park, CA 94025, USA*

(c) *Max-Planck-Institut für Kernphysik, D-69029, Heidelberg, Germany*

(d) *Institut für Synchrotronstrahlung, FZ Karlsruhe, D-76021 Karlsruhe, Germany*

(e) *Kirchhoff-Institut für Physik, Universität Heidelberg, D-69120 Heidelberg, Germany*

(f) *Fachbereich Physik, Universität Mainz, D-55099 Mainz, Germany*

(g) *CERN, CH-1211 Geneva 23, Switzerland*

(h) *Department of Physics, Princeton University, Princeton NJ 08544, USA*

(i) *Experimental Physik IV, Universität Dortmund, D-44221 Dortmund, Germany*

Presenter: N.O. Hashim (hashim@hep.physik.uni-siegen.de), ger-hashim-NO-abs1-ORAL

Cosmic muons are produced through interactions of primary cosmic radiation in the atmosphere. They are a component of extensive air showers which can also be measured underground. The CosmoALEPH experiment used the ALEPH detector at the European Laboratory for Particle Physics, CERN, to measure cosmic muon events at a depth of 320 mwe underground. The momentum spectrum and charge ratio of the cosmic muons are measured. The results are compared with the expectations from MC simulations based on different hadronic interaction models.

1. Introduction

Since their discovery, great progress has been achieved in the field of cosmic ray physics particularly towards the understanding of the origin, transport and acceleration mechanisms of the high energy particles that constitute primary cosmic rays, their interaction processes in the galactic and extra galactic media, and also in the Earth's atmosphere[1]. The interaction of primary cosmic ray particles in the Earth's atmosphere leads to production of a cascade of secondary particles or Extensive Air Showers (EAS) with various components - electromagnetic, hadronic, muon and neutrino components. There is a large number of models to describe these interactions.

Many cosmic ray experiments have used a variety of observables in EAS that provide an understanding of the hadronic interactions and also shed some light on the chemical composition of the primary particles. The muon flux at the surface provides a useful tool for the calculations of neutrino fluxes, the reconstruction of EAS and it can serve as a test of various interaction models.

The CosmoALEPH detector, which was one of the experiments in CosmoLEP[2, 3] used the ALEPH detector at the European Laboratory for Particle Physics, CERN, to measure the muonic component of EAS. Preliminary results[4] have recently shown that the momentum spectrum and charge ratio for cosmic muons measured by CosmoALEPH are well within the world average [5]. This work reports on further improvements in the reconstruction of the cosmic muon events and data analysis.

2. Experimental Setup

The ALEPH detector and its applicability as a tool for cosmic ray measurements has been well described [6, 7, 8, 9]. It was located at a depth of 320 mwe, which corresponds to an energy cut-off of about 70 GeV for

vertical muons. For measurements of cosmic muon events, the hadron calorimeter HCAL and time projection chamber TPC have been used. The trigger efficiency for HCAL is about 85% for vertical muons and a spatial resolution of about $160\mu\text{m}$ in the TPC allows the measurements of up to 3 TeV. A total of about 1.1 million cosmic muon events from dedicated cosmic runs are analysed in this work. The data are compared with Monte Carlo simulations.

3. Data Selection and Analysis

A total of 65,946 vertical cosmic muons, zenith angle θ up to 10° , with momentum greater than 5 GeV measured in ALEPH were selected for analysis. The muon energies at the surface E_μ are extrapolated from their respective energies measured in ALEPH E_N at the depth R using the relation,

$$E_\mu = \frac{a}{b} \left(e^{bR/\cos\theta} - 1 \right) + E_N \cdot e^{bR/\cos\theta}$$

which takes into account the energy losses due to ionisation, bremsstrahlung and direct electron pair production described by the terms a and b respectively[1]. The geometrical acceptance is corrected for the non-uniformity in the azimuth angle distribution using MC simulation. The data are then corrected for the track reconstruction efficiency which is estimated to be about 79%. The momentum spectrum is unfolded using Bayesian technique [10] taking into account the detector resolution and energy losses in the overburden. This is achieved by a complete simulation of cosmic muon events in the CosmoALEPH experiment using the following programs; CORSIKA[11] for air shower simulations, GEANT[12] for muon transport and interactions in ALEPH including the overburden. A detailed description is in preparation [13].

4. Results and Discussion

The preliminary momentum spectrum and the charge ratio for vertical muons at the surface is shown in Figure 1. The data are reasonably comparable with results of earlier experiments and a recent compilation [5]. The charge ratio, R_μ , of positive to negative muons obtained has an average value of 1.278 ± 0.011 . Only statistical uncertainties are shown. Systematic uncertainties are expected to be below 10%. A comparison with the Monte Carlo Simulations based on different hadronic interaction models is shown on Figure 2. The MC simulations are normalised to the selected number of events in the data. The slope of the number of muons at the surface predicted by the models agrees very well with our data. However, some systematic effects are evident in the calculations of the charge ratio. SIBYLL, which is known to generate a relatively lower muon content[14, 15], shows systematically higher values for the muon charge ratio compared to our measurements. The muon flux is closely related to the neutrino flux and the muon charge ratio is a sensitive observable for precision tests of muon and neutrino production in EAS [16].

5. Outlook

The preliminary results show a momentum spectrum that compares very well with previous measurements and the charge ratio is well described by Monte Carlo simulations. This work shows that the charge ratio for cosmic muons measured in CosmoALEPH is indeed an observable in EAS that can be used for testing theoretical hadronic interaction models. Further work is in progress to compare the data with updated models




Figure 1. Momentum spectrum and charge ratio for vertical muons compared with other experiments[5, 17].




Figure 2. Number of vertical muons at the surface and their charge ratio compared with MC simulations based on different hadronic interaction models.

for high energy interactions such as QGSJET-II [18] and NEXUS 3 [19] which will be included in the next version of CORSIKA[15, 20]. A more detailed analysis of the detector acceptance and trigger efficiency as function of muon momentum for inclined muons is in progress.

6. Acknowledgments

One of the authors (NOH) gratefully acknowledges the funding from the German Academic Exchange Service, DAAD. The authors gratefully acknowledge the help of the ALEPH collaboration, in particular the shift crew members of ALEPH, without whose support the data taking could not have been possible. The data analysis of the CosmoALEPH experiment has been funded by the Deutsche Forschungsgemeinschaft under grant DFG/Gr/1796/1-3.

References

- [1] C. Grupen. *Astroparticle Physics*. Springer-Verlag, Berlin Heidelberg, 2005. 441 p.
- [2] H. Wachsmuth, Proposal to search for cosmic ray coincidences in the LEP-Detectors, Draft, 16 December 1993, CosmoLEP-Note 94.000.
- [3] A. Ball, G. Bonvicini, C. Grupen et al., CERN/LEPC 94-10, LEPC/M, 1994, p 109
- [4] D. Zimmermann et al. The cosmic muon spectrum and charge ratio in CosmoALEPH. NIM A 525 (2004) pp 141 - 143.
- [5] T. Hebbeker and C. Timmermans. A compilation of high energy atmospheric muon data at sea level *Astropart. Phys.* 18 (2002) p 107-127.
- [6] ALEPH collaboration. Performance of the ALEPH detector at LEP. NIM A 360 (1995) pp 481-506.
- [7] M. Schmelling et al. Studying extendend air showers with CosmoALEPH. First Arctic Workshop on Cosmic Ray Muons, Sodankylae, Finland, April 24-29, 1999.
- [8] H. Besier et al. Cosmic ray physics with the ALEPH detector. NIM A 454 (2000) pp 201 - 206.
- [9] C. Grupen et al. Measurements of the muon component of extensive air showers at 320 m.w.e. underground. NIM A 510 (2003) pp 190-193
- [10] G. D'Agostini. A multidimensional unfolding method based on Bayes' theorem. NIM A362 (1995) pp 487-498.
- [11] D. Heck et al. CORSIKA: A Monte Carlo code to simulate extensive air showers. FZKA 6019, Forschungszentrum Karlsruhe (1998)
- [12] CERN. GEANT 3.21. Detector description and simulation tool. CERN Library Long Write-up W5013, CERN, GENEVA. 1994.
- [13] N.O. Hashim et al. Simulation of cosmic muon events in CosmoALEPH. In preparation.
- [14] J.R. Hörandel et al. Test of High Energy Hadronic Interaction Models using EAS Data. Proc. 26th Int. Cosmic Ray Conference, Salt Lake City 1 (1999) p 131 (and FZKA 6345, Forschungszentrum Karlsruhe (1999))
- [15] D. Heck. The influence of hadronic interaction models on simulated air showers: A phenomenologic comparison. Lecture Notes. VIHROS CORSIKA School 2005. May 29 - June 5 2005. Freudenstadt, Germany.
- [16] J. Wentz et al. The relevance of the muon charge ratio for the atmospheric neutrino anomaly. *J.Phys. G: Nucl. Part. Phys.* 27 (2001) pp 1699-1708.
- [17] The L3 Collaboration. Measurement of the atmospheric muon spectrum from 20 to 3000 GeV. CERN-PH-EP-2004-023. June 1, 2004.
- [18] S. Ostapchenko. Non-linear effects in high energy hadronic interactions. hep-ph/0501093 Jan. 2005.
- [19] K. Wener et al. A new string model: NEXUS 3. hep-ph/0306151 June 2003
- [20] D. Heck et al. Influence of low-energy hadronic interaction programs on air shower simulations with CORSIKA. Proc. 28th ICRC, Tsukuba, Japan, 1 (2003) p 279.