

## A Cosmic Ray Trigger for LOFAR: First results

S. Lafebre<sup>a</sup>, A. Nigl<sup>a</sup>, W.D. Apel<sup>b</sup>, F. Badea<sup>b</sup>, L. Bähren<sup>c</sup>, K. Bekk<sup>b</sup>, A. Bercuci<sup>d</sup>,  
M. Bertaina<sup>e</sup>, P.L. Biermann<sup>f</sup>, J. Blümer<sup>b,g</sup>, H. Bozdog<sup>b</sup>, I.M. Brancus<sup>d</sup>, S. Buitink<sup>a</sup>,  
M. Brüggemann<sup>h</sup>, P. Buchholz<sup>h</sup>, H. Butcher<sup>c</sup>, A. Chiavassa<sup>e</sup>, K. Daumiller<sup>b</sup>,  
A.G. de Bruyn<sup>c</sup>, C.M. de Vos<sup>c</sup>, F. Di Pierro<sup>e</sup>, P. Doll<sup>b</sup>, R. Engel<sup>b</sup>, H. Falcke<sup>a,c,f</sup>,  
H. Gemmeke<sup>i</sup>, P.L. Ghia<sup>j</sup>, R. Glasstetter<sup>k</sup>, C. Grupen<sup>h</sup>, A. Haungs<sup>b</sup>, D. Heck<sup>b</sup>,  
J.R. Hörandel<sup>g</sup>, A. Horneffer<sup>a,f</sup>, T. Huege<sup>b,f</sup>, K.-H. Kampert<sup>k</sup>, G.W. Kant<sup>c</sup>, U. Klein<sup>l</sup>,  
Y. Kolotaev<sup>h</sup>, Y. Koopman<sup>c</sup>, O. Krömer<sup>i</sup>, J. Kuijpers<sup>a</sup>, G. Maier<sup>b</sup>, H.J. Mathes<sup>b</sup>,  
H.J. Mayer<sup>b</sup>, J. Milke<sup>b</sup>, B. Mitrica<sup>d</sup>, C. Morello<sup>j</sup>, G. Navarra<sup>e</sup>, S. Nehls<sup>b</sup>, R. Obenland<sup>b</sup>,  
J. Oehlschläger<sup>b</sup>, S. Ostapchenko<sup>b</sup>, S. Over<sup>h</sup>, H.J. Pepping<sup>c</sup>, M. Petcu<sup>d</sup>, J. Petrovic<sup>a</sup>,  
T. Pierog<sup>b</sup>, S. Plewnia<sup>b</sup>, H. Rebel<sup>b</sup>, A. Risse<sup>m</sup>, M. Roth<sup>g</sup>, H. Schieler<sup>b</sup>, G. Schoonderbeek<sup>c</sup>,  
O. Sima<sup>d</sup>, M. Stümpert<sup>g</sup>, G. Toma<sup>d</sup>, G.C. Trinchero<sup>j</sup>, H. Ulrich<sup>b</sup>, S. Valchierotti<sup>e</sup>,  
J. van Buren<sup>b</sup>, W. van Capellen<sup>e</sup>, W. Walkowiak<sup>h</sup>, A. Weindl<sup>b</sup>, S. Wijnholds<sup>c</sup>, J. Wochele<sup>b</sup>,  
J. Zabierowski<sup>m</sup>, J.A. Zensus<sup>f</sup>, D. Zimmermann<sup>h</sup>

(a) Dept. of Astrophysics, IMAPP, Radboud University, 6525 ED Nijmegen, The Netherlands

(b) Institut für Kernphysik, Forschungszentrum Karlsruhe, 76021 Karlsruhe, Germany

(c) ASTRON, 7990 AA Dwingeloo, The Netherlands

(d) National Institute of Physics and Nuclear Engineering, 7690 Bucharest, Romania

(e) Dipartimento di Fisica Generale dell'Università, 10125 Torino, Italy

(f) Max-Planck-Institut für Radioastronomie, 53121 Bonn, Germany

(g) Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76021 Karlsruhe, Germany

(h) Fachbereich Physik, Universität Siegen, 57068 Siegen, Germany

(i) Inst. Prozessdatenverarbeitung und Elektronik, Forschungszentrum Karlsruhe, 76021 Karlsruhe, Germany

(j) Istituto di Fisica dello Spazio Interplanetario, INAF, 10133 Torino, Italy

(k) Fachbereich Physik, Universität Wuppertal, 42097 Wuppertal, Germany

(l) Radioastronomisches Institut der Universität Bonn, 53010 Bonn, Germany

(m) Soltan Institute for Nuclear Studies, 90950 Lodz, Poland

Presenter: Sven Lafebre (s.lafebre@astro.ru.nl), net-lafebre-S-abs1-he15-oral

We present the first results of an independent cosmic ray trigger for the multiple dipole-antenna radiotelescope LOFAR (LOw Frequency ARray). LOPES (the LOfar PrototypE Station), at the KASCADE (KARlsruhe Shower Core and Array DETector) site in Germany has been initiated as a test case for LOFAR, and designed to detect air showers through coherent radiation pulses from air showers upon external triggers by particle detectors. To fully exploit the capabilities in detecting CRs with the final LOFAR telescope, however, an independent, radio-only trigger is needed, which has been developed from experience with LOPES and other LOFAR test stations. Here, we present the first results of the application of such a trigger, and discuss optimization of the different parameters.

The material covered in this talk overlaps with what is to be presented in a poster by A. Nigl, *A Cosmic Ray Trigger for LOFAR*, **net-nigl-abs2-he15-poster**. Please refer to this paper.