



Realistic modelling of the termination shock position using Voyager 2 solar wind data

J.L.SNYMAN¹, S.E.S.FERREIRA¹, M.S.POTGIETER¹ AND K.SCHERER²

¹Unit for Space Physics, School of Physics, North-West University, 2520 Potchefstroom, South Africa

²Institut für Theoretische Physik, Lehrstuhl IV: Weltraum- und Astrophysik, Ruhr-Universität Bochum, D-44780 Bochum, Germany

20131232@nwu.ac.za

Abstract: The termination shock position in time is calculated from Voyager 2 solar wind data, using a multi species hydrodynamic model, also including the latitudinal variations in the solar wind velocity and density due to the 11-year solar cycle. The results are compared to those obtained by Webber (2005) and the possibility of Voyager 2 crossing the TS in termination shock in the near future is investigated.

Introduction

Although the Voyager 1 spacecraft crossed the solar wind (SW) termination shock (TS) in December 2004, its approach to the TS is still a topic of interest. Specifically, in [11] the < 5 MeV proton intensities measured by Voyager 1 from 2002.6 onward is explained as a result of Voyager 1 coming close to the TS in 2002. After 2002 the TS moved outward with a velocity comparable to that of Voyager 1, causing the spacecraft to have several close encounters with the shock before finally crossing it in December 2004.

The position of the TS with time is calculated in [11] using both long and short term SW pressure variations observed at Voyager 2. In [11] the Voyager 2 SW data is extrapolated back to 1 AU and used to calculate the subsequent TS response using results from a range of models, notably [3, 5, 9, 10, 12] and [13]. The approach followed by [11] is empirical, and 'response coefficients' are defined in correlation with results from the studies cited above.

In this paper, we would like to add to the result from [11] in the following ways: in extrapolating the Voyager 2 SW data back to 1 AU, [11] assumes a r^{-2} dependence for the SW density (where r denotes distance from the Sun) as well as a constant

SW velocity. The former assumption is almost exact, even in a heliosphere where the multi-species interaction between the SW and the local interstellar medium (LISM) are considered. However, we extend the study by assuming that the SW velocity decreases with increasing radial distance due to the interaction between the SW and neutral hydrogen producing pick-up ions (PUI's). Furthermore we use the extrapolated values as SW boundary conditions in a hydrodynamic model of the heliosphere, which includes the mutual effects of protons, neutral hydrogen and PUI's self consistently together with the 11-year solar cycle. Results are calculated in a plane parallel to the solar rotation axis, which allows for a realistic inclusion of latitudinal effects, such as the appearance and disappearance of the fast SW.

Model and data

The hydrodynamical model used in this study is derived from [1], where the Euler equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0 \quad (1)$$

$$\frac{\partial}{\partial t} (\rho \mathbf{u}) + \nabla \cdot (\rho \mathbf{u} \otimes \mathbf{u}) + \nabla P = 0 \quad (2)$$

$$\frac{\partial E}{\partial t} + \nabla \cdot \left(\mathbf{u} \left(\frac{P}{\gamma - 1} + \frac{1}{2} \rho u^2 \right) \right) = 0 \quad (3)$$

are solved for each different particle species, on a plane-polar grid, corresponding to a plane parallel to the solar axis of rotation. The model takes into account the mutual effect of protons, hydrogen and PUI's. The model was adapted to include the latitudinal variation of both the SW density and velocity (see [2] and [8]) over the 11-year solar cycle.

From the model results (see [1]) the SW density is almost exactly proportional to r^{-2} , while the SW velocity decreases almost linearly with increasing radial distance. Using

$$v_p = mr + c \quad (4)$$

to approximate the decrease of the SW velocity v_p with r , and assuming that the SW decreases to 5% of its original value, the values of m and c are

$$m = -1.69 \times 10^{-9} \text{ s}^{-1} \quad c = 401.26 \text{ km.s}^{-1} \quad (5)$$

It should be noted that the choice of 5% depends on the assumed SW density at 1 AU as well as the assumed LISM hydrogen density outside the heliosphere. The choice of 5% is fairly accurate for SW densities between 5 cm^{-3} and 7 cm^{-3} , with a LISM hydrogen density assumed to be 0.1 cm^{-3} . The time Δt it took for a fluid particle to travel a distance r and reach a velocity v_p is calculated to be

$$\Delta t = \frac{1}{m} \ln \left| \frac{mr}{x} + 1 \right|. \quad (6)$$

Using Eq.(4), the SW velocity observed at a certain radial position r and time t is extrapolated back to a velocity at 1 AU. From Eq.(6) the corresponding time value is calculated as $t - \Delta t$. In using this procedure, the daily averaged Voyager 2 SW data is extrapolated back to 1 AU and the corresponding 26-day average values are calculated and shown in Figure 1.

The 26-day averaged values from Figure 1 are used as an inner boundary condition to the hydrodynamic model. During solar maximum it is as-




Figure 1: 26-day averages of Voyager 2 solar wind (SW) data, extrapolated back to 1 AU. Original data taken from <http://cohoweb.gsfc.nasa.gov>

sumed that the Voyager 2 data extends over all latitudes. As the model progresses to solar minimum, the influence of the Voyager 2 data is gradually decreased at higher latitudes, so that during solar minimum the density and velocity over high latitudes are smooth. This corresponds to observations made by Ulysses, indicating less solar wind variability at high latitudes (see high latitude Ulysses data from <http://cohoweb.gsfc.nasa.gov> for example).

Regarding the crossing of the TS by Voyager 1, we present two cases. Firstly, following [11], it is assumed that Voyager 1 came close to the TS in 2002.6. For the purpose of this work we choose the distance between Voyager 1 and the TS at 2002.6 as 1 AU. Secondly, regarding the consensus that Voyager 1 crossed the TS in 2004.9, we present a second case where it is assumed that Voyager 1 encountered the shock in 2004.9. Results are presented in the next section.




Figure 2: The termination shock (TS) position as it varies with time, where it is assumed that Voyager 1 was 1 AU away from the TS in 2002.6. The solid black line shows the TS position along Voyager 1's trajectory, while the grey line shows the TS position along the trajectory of Voyager 2. The dotted vertical lines correspond to 2002.6 and 2004.9 respectively, while the horizontal dotted line corresponds to 94 AU.

Results

The position of the termination shock r_{TS} is calculated as it varies in time due to the SW fluctuations measured by Voyager 2. Following [11], the distance between Voyager 1 and the TS is assumed to be 1 AU in 2002.6. The result is shown in Figure 2. It was found in [11] that Voyager 1 had a series of close encounters with the TS between 2002 and 2004, which agrees with Figure 2. However, the results imply that Voyager 1 moved clear of the TS as early as 2003.8, which is difficult to reconcile with the consensus that the crossing occurred in 2004.9. Furthermore, the results also imply that Voyager 2 should have already crossed the TS in 2005.5.

Regarding the second case, it is assumed that Voyager 1 crossed the shock at 2004.9. The results are shown in Figure 3. The results show that Voyager 1 and the TS had comparable outward velocities between 2002 and 2004 (which agrees with [11]), whereafter the TS moved inward, resulting in Voyager 1 crossing the TS in 2004.9. Furthermore, the projected motion of the TS implies that Voyager 2 will probably encounter the TS as early as 2007.5.

Regarding the above mentioned results, it needs to be noted that the further Voyager 1 was from the




Figure 3: The termination shock (TS) position as it varies with time, where it is assumed that Voyager 1 crossed the TS in 2004.9. The solid black line shows the TS position along Voyager 1's trajectory, while the grey line shows the TS position along the trajectory of Voyager 2. The dotted vertical lines correspond to 2002.6 and 2004.9 respectively, while the horizontal dotted line corresponds to 94 AU.

shock between 2002 and 2004, the more difficult it becomes to explain the > 0.5 MeV proton data (see [11]) observed by Voyager 1 during this period. In order to constrain the TS position relative to both the Voyager spacecraft trajectories accurately, it needs to be established how far away from the TS increases in < 0.5 MeV protons, as detected by Voyager 1, will be observed.

Lastly it should be noted that neither the effects of the heliospheric magnetic field (HMF) nor that of the inter stellar magnetic field (IMF) are included in the model. The inclusion of the HMF and IMF can alter the geometry of the heliosphere, which in turn will have an effect on the predicted time at which Voyager 2 will cross the shock as well as charged particle intensities seen by both spacecraft close to the TS (see [4, 6] and [7]).

Summary and conclusion

Using Voyager 2 SW observations to calculate the time dependent position of the TS indicates that, as Voyager 1 approached the TS, the TS itself was moving outward with a velocity comparable to that of Voyager 1. This resulted in Voyager 1 moving close to the shock for a prolonged period of time (between 2002 and 2004) before finally crossing

the shock. Exactly how close still needs to be constrained by investigating < 0.5 MeV proton intensities close to the shock. It is unlikely that Voyager 1 came as close to 1 AU in 2002.6 since this would imply that Voyager 2 already crossed the shock as early as 2005.5. In assuming that Voyager 1 crossed the TS in 2004.9, it is found that Voyager 2 is likely to cross the TS as early as 2007.5.

Acknowledgements

The financial assistance of the National Research Foundation (NRF) towards this research is hereby acknowledged. Opinions expressed and conclusions arrived at, are those of the author and are not necessarily to be attributed to the NRF

References

- [1] H. J. Fahr, T. Kausch, and H. Scherer. A 5-fluid hydrodynamic approach to model the solar system-interstellar medium interaction. , 357:268–282, May 2000.
- [2] S. E. S. Ferreira and K. Scherer. Time Evolution of Galactic and Anomalous Cosmic-Ray Spectra in a Dynamic Heliosphere. *Astrophys. J.*, 642:1256–1266, May 2006.
- [3] V. Izmodenov, G. Gloeckler, and Y. Malama. When will Voyager 1 and 2 cross the termination shock? , 30:3–1, April 2003.
- [4] V. V. Izmodenov and D. B. Alexashov. Multi-component 3D modeling of the heliospheric interface: effects of interstellar magnetic field. In J. Heerikhuisen, V. Florinski, G. P. Zank, and N. V. Pogorelov, editors, *Physics of the Inner Heliosheath*, volume 858 of *American Institute of Physics Conference Series*, pages 14–19, September 2006.
- [5] J. Y. Lu, Y. C. Whang, and L. F. Burlaga. Interaction of a strong interplanetary shock with the termination shock. , 104:28249–28254, December 1999.
- [6] M. Opher, E. C. Stone, and P. C. Liewer. The Effects of a Local Interstellar Magnetic Field on Voyager 1 and 2 Observations. , 640:L71–L74, March 2006.
- [7] N. V. Pogorelov, G. P. Zank, and T. Ogino. Three-dimensional Features of the Outer Heliosphere due to Coupling between the Interstellar and Interplanetary Magnetic Fields. II. The Presence of Neutral Hydrogen Atoms. , 644:1299–1316, June 2006.
- [8] K. Scherer and S. E. S. Ferreira. A heliospheric hybrid model: hydrodynamic plasma flow and kinetic cosmic ray transport. *Astrophysics and Space Sciences Transactions*, 1:17–27, March 2005.
- [9] T. R. Story and G. P. Zank. Response of the termination shock to interplanetary disturbances 2. MHD. , 102:17381–17394, August 1997.
- [10] C. Wang and J. W. Belcher. The heliospheric boundary response to large-scale solar wind fluctuations: A gasdynamic model with pickup ions. , 104:549–556, January 1999.
- [11] W. R. Webber. An empirical estimate of the heliospheric termination shock location with time with application to the intensity increases of MeV protons seen at Voyager 1 in 2002-2005. *Journal of Geophysical Research (Space Physics)*, 110:10103–10109, October 2005.
- [12] Y. C. Whang and L. F. Burlanga. Termination shock - Solar cycle variations of location and speed. , 98:15221–15230, September 1993.
- [13] G. P. Zank and H.-R. Müller. The dynamical heliosphere. *Journal of Geophysical Research (Space Physics)*, 108:7–1, June 2003.