



## The effect of a modified Parker field on the modulation of the galactic cosmic rays

SHOKO MIYAKE, SHOHEI YANAGITA.

Faculty of Science, Ibaraki University, Mito 310-8512, Japan

06nd408g@mcs.ibaraki.ac.jp

**Abstract:** The effects on the galactic cosmic ray modulation of a Parker field modified by the motion of footpoints of the magnetic field lines and the latitudinal dependent solar wind velocity are numerically investigated. The calculation of the propagation of the galactic cosmic ray is made by solving a coupled set of stochastic differential equations (SDE) which is equivalent to the so-called diffusion convection partial differential equation. The three-dimensional code based on the SDE method has been developed into the modified Parker field. We present the details of the effects of latitudinal footpoints motion on the solar modulation of the galactic cosmic rays.

### Introduction

The heliospheric magnetic field (HMF) plays an important part in the model for the solar modulation of the galactic cosmic rays. The model for HMF by Parker [6] has been successfully used in modulation studies, however, this model may be an oversimplification to describe the field in the polar region of the heliosphere. Recently some models of HMF have been proposed to take account of the footpoint motion of open magnetic fields (e.g. [3], [4]). When we adopt the footpoint velocity  $u_{\perp} = R_{ss}\omega_{\theta}\hat{e}_{\theta} + R_{ss}\omega_{\phi}\hat{e}_{\phi}$  on the source surface in a frame rotating with the Sun, the HMF takes the following form

$$\begin{aligned} \mathbf{B} &= B_{ss} \left( \frac{R_{ss}}{r} \right)^2 \\ &\times \left( \hat{e}_r - \frac{\omega_{\theta} r}{V} \hat{e}_{\theta} - \frac{r(\Omega_{\odot} - \omega_{\phi}) \sin\theta}{V} \hat{e}_{\phi} \right) \\ &\times [1 - 2H(\theta - \theta')] , \end{aligned} \quad (1)$$

where  $r$  is the heliocentric distance,  $\theta$  is the co-latitude,  $R_{ss}$  is the radius of the source surface,  $B_{ss}$  is the magnetic field strength at the source surface, and  $\Omega_{\odot}$  is the rotational angular velocity of the Sun,  $V$  is the solar wind velocity,  $H$  is the Heaviside function, and  $\theta'$  is the co-latitude of the heliospheric current sheet. Note that this equation can be applicable only for a constant solar wind velocity  $V$ . The solar wind velocity has a latitudinal

dependence, especially near at the solar minimum phase. Schwadron and McComas [7] studied the effect of the latitudinal dependent solar wind on the HMF (eq.1) and the cosmic ray acceleration at the termination shock. Burger and Sello [2] calculated the latitudinal gradient of the galactic cosmic rays by adopting a simple two-dimensional HMF which takes account of the latitudinal dependent solar wind velocity. The direction of the latitudinal footpoint motion affects the structure of the HMF in the region where the solar wind velocity changes significantly with latitude. We have numerically investigated the solar modulation of the galactic cosmic rays by the same HMF assumed by Burger and Sello [2]. We present details of the effects of the direction of the latitudinal footpoint motion on the solar modulation of the galactic cosmic rays.

### Numerical models

Transport of the galactic cosmic rays in the heliosphere is described by the so-called diffusion convection partial differential equation. This diffusion convection equation is equivalent to a coupled set of stochastic differential equations (SDE) [8, 9]. This SDE method allows us to get some information about the solar modulation phenomena of the galactic cosmic ray not obtained by the

other numerical methods, such as the distributions of arrival time, energy lost, and trajectory. The stochastic numerical code adapted for the wavy heliospheric current sheet in the standard Parker field has been developed by Miyake and Yanagita [5]. We have developed this three-dimensional code based on the SDE method into a Parker field modified by the latitudinal footpoint motion and the latitudinal dependent solar wind velocity. The SDE equivalent to the diffusion convection partial differential equation is written as

$$d\mathbf{X} = (\nabla \cdot \boldsymbol{\kappa} - \mathbf{V}(\theta) - \mathbf{V}_d) dt + \sum_s \boldsymbol{\sigma}_s dW_s(t), \quad (2)$$

$$dP = \frac{1}{3} P (\nabla \cdot \mathbf{V}(\theta)) dt,$$

where  $\mathbf{X}$  and  $P$  are the position and the momentum of the particle,  $\mathbf{V}(\theta)$  is the solar wind velocity which has a latitudinal dependence,  $\mathbf{V}_d$  is the gradient-curvature drift velocity,  $\boldsymbol{\kappa}$  is the diffusion coefficient tensor,  $\sum_s \boldsymbol{\sigma}_s^\mu \boldsymbol{\sigma}_s^\nu = 2\kappa^{\mu\nu}$ , and  $dW_s$  is a Wiener process given by the Gaussian distribution. We adopted  $\kappa_{\parallel} = 1.5 \times 10^{21} \beta(p/(1\text{GeV}/c))(Be/B) \text{ cm}^2/\text{s}$  and  $\kappa_{\perp} = 0.05\kappa_{\parallel}$ . We have used the ‘‘drift velocity field method’’ [1] for the calculation of drift in the heliospheric current sheet. We investigated a case of the tilt angle of the heliospheric current sheet  $\alpha$  fixed at  $10^\circ$ .

In our simulation, particles start at 1 AU on the equatorial plane and run backward in time until they exit the heliospheric boundary, 90 AU. The momentum spectrum  $f_{\mathbf{X}}(p)$  at arbitrary position  $\mathbf{X}$  is written as a convolution of the spectrum  $f_{\mathbf{X}_0}(p_0)$  at the heliospheric boundary with the normalized transition probability  $F(p_0, \mathbf{X}_0|p, \mathbf{X})$  obtained by our SDE method as

$$f_{\mathbf{X}}(p) = \int f_{\mathbf{X}_0}(p_0) F(p_0, \mathbf{X}_0|p, \mathbf{X}) dp_0. \quad (3)$$

Here the spectrum at the boundary is  $f_{\mathbf{X}_0}(p_0) \propto (m^2 c^4 + p_0^2 c^2)^{-1.85}/p_0$  which is assumed to be uniform at the heliospheric boundary.

We adopted the simplified model of HMF which is assumed by Burger and Sello [2] and takes account of the footpoint motion of open magnetic fields and




Figure 1: The solar wind velocity profile used in this study.




Figure 2: (a) The effect of the solar wind velocity profile shown in Figure 1 on the HMF strength at 1 AU where  $r_p \times dV/d\theta \leq 0$ ; (b) Same as (a) where  $r_p \times dV/d\theta \geq 0$ .

the latitudinal dependent solar wind velocity,

$$\mathbf{B} = B_{ss} \left( \frac{R_{ss}}{r} \right)^2 \times \left[ \left( 1 - \frac{r_p}{4V(\theta)} \frac{dV(\theta)}{d\theta} \right) \hat{\mathbf{e}}_r - \frac{r\Omega_{\odot} \sin\theta}{V(\theta)} \hat{\mathbf{e}}_{\phi} \right], \quad (4)$$

where  $r_p$  is a dimensionless constant. Figure 1 shows the solar wind velocity profile used in this study. We assumed the solar wind velocity depends only on  $\theta$ . Figure 2.a shows the effect of the solar wind velocity profile shown in Figure 1 on the HMF strength at 1 AU where  $r_p \times dV/d\theta \leq 0$ . The solid line indicates the HMF strength where the absolute value of  $r_p$  is 4. The dotted line indicates the HMF strength where  $r_p = 0$ , namely for the standard Parker field. Figure 2.b is the same as Figure 2.a for  $r_p \times dV/d\theta \geq 0$ . While the difference of the HMF from the standard Parker model




Figure 3: The latitudinal dependence of the 1 GeV proton intensities at 1 AU where  $r_p \times dV/d\theta \leq 0$ . The intensity is normalized to the intensity at the heliospheric boundary. The solid line indicates the intensities where the absolute value of  $r_p$  is 4. The dotted line indicates the intensities where  $r_p = 0$ .

for both cases is localized in the region where the solar wind velocity changes significantly with latitude, the feature of the difference is quite different qualitatively between Fig 2.a and b. These differences would affect the solar modulation of the galactic cosmic rays. We have numerically investigated the effect of these differences on the modulation of the galactic cosmic rays.

## Results and Conclusions

Figure 3 shows the latitudinal dependence of the 1 GeV proton intensities at 1 AU where  $r_p \times dV/d\theta \leq 0$  for both signs of the magnetic polarity  $qA$ . The intensity is normalized to the intensity at the heliospheric boundary. The solid line indicates the intensities where the absolute value of  $r_p$  is 4. The dotted line indicates the intensities where  $r_p = 0$ , namely for the standard Parker field. Figure 4 is the same as Figure 3 for  $r_p \times dV/d\theta \geq 0$ . In Figure 3, the intensities for the modified Parker field where  $|r_p| = 4$  is lower than the intensities for the standard Parker field, especially in the region where the solar wind velocity changes significantly with latitude. This tendency comes from




Figure 4: Same as Figure 3 but  $r_p \times dV/d\theta \geq 0$ .

the dependence of the diffusion mean free path and the drift scale on the HMF strength. In our simulation, the diffusion coefficient and the drift scale are inversely proportional to the HMF strength. Accordingly it is difficult for the galactic cosmic rays to access to the region where the HMF strength is relatively high, i.e. in the region where the solar wind velocity changes significantly with latitude. As seen in Figure 3, this tendency is opposite to the results by Burger and Sello [2]. This contradiction in the results between Burger and Sello [2] and ours may come from the difference in the assumption of the dependence of the diffusion coefficient on the HMF strength. Burger and Sello [2] adopted a model the parallel mean free path of the diffusion is proportional to  $B^{5/3}$  at low rigidities and is independent of  $B$  at high rigidities. Therefore the different latitudinal dependence may result in for the case of  $r_p \times dV/d\theta \leq 0$ . Fig 4 shows the same results shown in Fig 3 for the case of  $r_p \times dV/d\theta \geq 0$ . The intensities in the region where the HMF strength is modified differ from those predicted for the case of the standard Parker model including the direction of the change.

Although the local drift patterns change near at the region where the solar wind velocity changes significantly with latitude, the global drift pattern has the same tendency as the pattern for the standard Parker field. For the positive polarity  $qA > 0$ , the galactic cosmic rays drift from the polar region towards the heliospheric current sheet, and

outward along the heliospheric current sheet. In contrast, for the negative polarity  $qA < 0$ , the direction of the drift is opposite to the case  $qA > 0$ . This is the reason for the decrease (increase) in the intensities for the modified Parker field at higher co-latitude than the region where the solar wind velocity changes significantly with latitude for  $qA > 0$  and at lower co-latitude for  $qA < 0$  when  $r_p \times dV/d\theta \leq 0$  ( $\geq 0$ ).

## Acknowledgements

Numerical computations were carried out on general common use computer system at the Center for Computational Astrophysics, CfCA, of the National Astronomical Observatory of Japan. S. Miyake is supported by a JSPS Research Fellowship.

## References

- [1] R. A. Burger, H. Moraal, and M. S. Potgieter. On the inclusion of a wavy neutral sheet in two-dimensional drift models. *Proc. 20th Int. Cosmic-Ray Conf. (Moscow)*, 3:283–286, 1987.
- [2] R. A. Burger and P. C. Sello. The effect on cosmic ray modulation of a Parker field modified by a latitudinal-dependent solar wind speed. *Adv. Space Res.*, 35:643–646, 2005.
- [3] L. A. Fisk. Motion of the footpoints of heliospheric magnetic field lines at the Sun: Implications for recurrent energetic particle events at high heliographic latitudes. *J. Geophys. Res.*, 101(A7):15547–15553, 1996.
- [4] J. R. Jokipii and J. Kóta. The polar heliospheric magnetic field. *Geophys. Res. Lett.*, 16(1):1–4, 1989.
- [5] S. Miyake and S. Yanagita. Effects of the tilted and wavy current sheet on the solar modulation of galactic cosmic rays. *Proc. 29th Int. Cosmic-Ray Conf. (Pune)*, 2:203–206, 2005.
- [6] E. N. Parker. Dynamics of the interplanetary gas and magnetic fields. *Astrophys. J.*, 128:664–676, 1958.
- [7] N. A. Schwadron and D. J. McComas. Heliospheric "FALTS": favored acceleration locations at the termination shock. *Geophys. Res. Lett.*, 30(11):1587, 2003.
- [8] Y. Yamada, S. Yanagita, and T. Yoshida. A stochastic view of the solar modulation phenomena of cosmic rays. *Geophys. Res. Lett.*, 25(13):2353–2356, 1998.
- [9] M. Zhang. A Markov stochastic process theory of cosmic-ray modulation. *Astrophys. J.*, 513:409–420, 1999.