



Particle acceleration in relativistic subluminal shock environments

A. MELI^a, J. K. BECKER^b, J. J. QUENBY^c AND J. LÜNEMANN^b

(a) *Department of Physics, University of Athens, 15783 Zografos, Athens Greece*

(b) *Institut für Physik, Universität, 44221 Dortmund Germany*

(c) *Imperial College London, Astrophysics Group, Blackett Laboratory SW7, 2BW, London UK*

jan.luenemann@uni-dortmund.de

Abstract: The understanding of the particle spectra resulting from acceleration in relativistic shocks as they occur in extragalactic sources is essential for the interpretation of the Cosmic Ray spectrum above the ankle ($E_p > 3 \cdot 10^{18}$ eV). It is believed that extragalactic sources like Active Galactic Nuclei and Gamma Ray Bursts can produce particle spectra up to $E_p \sim 10^{21}$ eV. In this contribution, subluminal shocks are investigated with respect to different shock boost factors Γ and the inclination angle between the shock normal and the magnetic field ψ . A correlation between the boost factor and the spectral behavior of the emitted particles is found. The results are compared to Gamma Ray Burst observations and the Cosmic Ray spectrum at the highest energies.

Introduction

The observations of electron synchrotron radiation in the radio regime and the X-ray variations from astrophysical environments such as Active Galactic Nuclei (AGN) and Gamma Ray Bursts (GRBs) indicate that there is a mechanism with which primary particles (protons or electrons) gain large amounts of energy via acceleration. Connected to this implied mechanism it is accepted that shocks are the source of these Cosmic Rays. The work of the late 70s by a number of authors such as [12, 3, 4, 1, 5] established the basic mechanism of particle 'diffusive' acceleration in non-relativistic shocks. Later on studies on relativistic shocks appeared (e.g. [17, 11, 19, 7]) and questioned the nature and the limits of the relativistic diffusive acceleration versus the non-relativistic shock mechanism. In this work, we present results on relativistic particle acceleration which are extended for relativistic source models (i.e. AGN, GRBs). These involve subluminal shocks with plasma shock velocity ranging from $\Gamma = 1 \rightarrow 1000$ Lorentz factor and both pitch angle and magnetic scattering. In the first section, the Monte Carlo simulations are described, while in the second section, the results are presented for different boost factors. In

the third section, the Monte Carlo spectra are used to calculate a possible contribution of AGN and GRBs to the observed spectrum of charged Cosmic Rays. The final section summarizes the results.

Numerical simulations

The Monte Carlo codes simulate relativistic oblique shocks with application to relativistic environments of AGN and GRBs. Initially the particles are considered relativistic, isotropically injected upstream of the shock and left to move towards the shock and along the way they collide with the scattering centers. Consequently, as they keep scattering between the upstream and downstream regions of the shock (its width is much smaller than the particle's gyroradius), they gain a considerable amount of energy. During our simulations a relativistic transformation is performed from the local plasma frames to the shock frame to check for shock crossings. The de Hofmann-Teller frame is used appropriately while the adiabatic invariant is conserved.

We change units so that $m \approx c = 1$. Particles leave the system if 'escaping' far downstream at the spatial boundary, or if reaching a defined maxi-

mum energy E_{max} for computational convenience, even though other physics describing particle escape or energy loss would probably need to be taken into account in realistic situations (forthcoming paper). The downstream spatial boundary required can be estimated from the solution of the convection-diffusion equation in a non-relativistic, large-angle scattering approximation in the downstream plasma. This gives the chance of return to the shock, $\exp(-V_2 r_b/d)$, yielding a probability of return of $2 \cdot 10^{-3}$.

The mean free path is calculated in the respective fluid frames by the formula: $\lambda = \lambda_c p_{1,2}$, assuming a momentum dependence to this mean free path for scattering along the field related to the spatial diffusion coefficient, κ , in the shock normal or x direction by $\kappa_{\parallel} = \lambda v/3$ and $\kappa = \kappa_{\parallel} \cos^2 \psi$ (for the oblique shocks). At the scattering centers the energy (momentum) of the particle is kept constant and only the direction of the velocity vector v is randomized, using a computational random number generator. Runs are performed with different spatial boundaries to investigate the effect of the size of the acceleration region on the spectrum, as well as to find a region where the spectrum is size independent. In the small-angle scatter case, the inherent anisotropy due to the high downstream sweeping effect may greatly modify this analytical estimate. A splitting technique is used similar to one used in Monte Carlo simulations of [15], [16], so that when an energy is reached such that only a few accelerated Cosmic Rays remain, each particle is replaced by a number of N particles of statistical weight $1/N$ so as to keep roughly the same number of Cosmic Rays being followed. Workers on the field [20] have demonstrated that small angle scattering (pitch angle diffusion) with $\delta\theta < 1/\Gamma$ applies with θ measured in the upstream fluid frame for scattering in a uniform field or a randomly oriented set of uniform field cells. This arises because particles attempting to penetrate upstream from the shock are swept back into the shock before they can move far in pitch. For this case, we allow particles' pitch angle diffusion to angles chosen at random, up to an angle $\sim 1/\Gamma$, where Γ is the upstream gamma, measured in the shock frame.

For the oblique shock case we apply here, provided the field directions encountered are reasonably isotropic in the shock frame, we know that

$\tan \psi_1 = \Gamma_1^{-1} \tan \psi_s \sim \Gamma_1^{-1} \sim \psi_1$ where '1' and 's' refer to the upstream and shock frames respectively. Given the current interest in following shock acceleration in the relativistic flow in test particle regime we outline a typical approach to such computations. Using the guiding center approximation, a test particle moving a distance, d , along a field line at ψ to the shock normal, in the plasma frame has a probability of collision within d given by $P(d) = 1 - \exp(-d/\lambda) = R$. Where the random number R is $0 \leq R \leq 1$. Weighting the probability by the current in the field direction, μ , then $d = -\lambda \mu \ln R$. Between the ' i th' and ' $i+1$ th' scattering shock crossing is checked by finding the position and time in the shock frame with $c = 1$ units, $\Delta x^{i+1} = \Gamma(\Delta x^i + V|\frac{\Delta x^i \sec \psi}{v\mu}|)$ and $\Delta t^{i+1} = \Gamma(|\frac{\Delta x^i \sec \psi}{v\mu}| + V\Delta x)$. We note that for the description of the physical quantities necessary throughout the simulations we use the fluid rest frames, the normal shock frame and the de Hoffman-Teller frame. Furthermore as the theory for the oblique shocks postulates [9], from the conservation of the first adiabatic invariant we can find the new pitch angle in the downstream frame and similar transformations allow the particle scattering to be followed in this frame.

Results

Energy spectra

Monte Carlo calculations as discussed above have been performed for different parameter settings. Three different shock obliquities have been simulated, i. e. $\psi = 23^\circ, 33^\circ$ and 43° , where ψ is the angle between shock normal and magnetic field. For each of the angles, nine different boost factors have been used, $\Gamma = 10, 20, 30, 100, 300, 500, 700, 900$ and 1000. The results will be presented in detail in [14]. Each single spectrum is then fit by a single power-law approximation, $dN/dE \propto E^{-\alpha_p}$, in order to get an estimate of the spectral index α_p . At the lowest energies, particle injection features are excluded. Since AGN and GRBs are likely to accelerate particles up to $\sim 10^{21}$ eV, this value is used as the maximum energy. This way, each shock boost factor Γ is associated with a spectral




Figure 1: Correlation between Γ and the spectral index for $\psi = 43^\circ$. The correlation appears to be linear.

index $\alpha_p(\Gamma)$. A linear correlation between α_p and Γ is found as it is displayed in Fig. 1. The figure shows the results for $\psi = 43^\circ$, the results for $\psi = 23^\circ$ and $\psi = 33^\circ$ are comparable. A linear fit is applied with a numerical result which is comparable for all three angles,

$$\alpha_p(\psi = 23^\circ) = 2.1 - 0.7 \cdot 10^{-3} \cdot \Gamma \quad (1)$$

$$\alpha_p(\psi = 33^\circ) = 2.0 - 0.9 \cdot 10^{-3} \cdot \Gamma \quad (2)$$

$$\alpha_p(\psi = 43^\circ) = 2.2 - 0.9 \cdot 10^{-3} \cdot \Gamma. \quad (3)$$

This implies three main things for relativistic, subluminal shocks:

1. The result is basically independent of the inclination angle between shock normal and magnetic field, as long as the shocks are oblique.
2. Mildly-relativistic shocks have spectral indices of around $\alpha_p \sim 2.0 - 2.4$.
3. The more relativistic the shock, the flatter the spectrum. Highly-relativistic shocks with $\Gamma \sim 1000$ can produce spectra as flat as $\alpha_p \sim 1.5$.

The above findings seem to support many observational evidences regarding irregular and flat spectra from many GRBs (further details in [14]).

Contribution to the charged Cosmic Ray spectrum

In this section, the spectra of different boost factors are associated to different source classes in order to calculate the possible contribution to the diffuse spectrum of charged Cosmic Rays. It is assumed in the following, that the Cosmic Ray spectrum above the ankle, $E > 10^{18.5}$ eV, is produced by protons from AGN and GRBs. For AGN, a fix boost factor of $\Gamma = 10$ is used. In the case of GRBs, it is assumed that 10% of all GRBs have $\Gamma = 100$, 80% have $\Gamma = 300$ and the remaining 10% have $\Gamma = 1000$. It is further assumed that the observed spectrum of charged Cosmic Rays is produced by a linear combination of AGN, dN_{AGN}/dE , and GRBs, dN_{GRB}/dE ,

$$\frac{dN_{CR}}{dE} = \alpha \cdot \frac{dN_{AGN}}{dE} + (1 - \alpha) \cdot \frac{dN_{GRB}}{dE}, \quad (4)$$

with $0 \leq \alpha \leq 1$. Source evolution has been taken into account by using the source density given by the observation of X-ray AGN [8]. This source evolution is also used in the case of GRBs, assuming that GRBs also follow the Star Formation Rate. The energy density of the observed Cosmic Ray spectrum j_E is then used to normalize the calculated Monte Carlo spectra

$$j_E := \int_{E_{\min}=10^{18.5} \text{ eV}} \frac{dN}{dE} E dE \quad (5)$$

$$\approx 10^{-7} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}. \quad (6)$$

The final result is shown in Fig. 2. The upper line represents a pure AGN spectrum, $\alpha = 1$, the middle line represents a mixture of 50% AGN and 50% GRBs ($\alpha = 0.5$) and the lower line is a pure GRB spectrum, $\alpha = 0$. Since the Cosmic Ray spectrum above $E = 10^{18.5}$ eV needs to be explained by extragalactic sources, the best fit results from a pure AGN spectrum. With a significant GRB contribution, the spectrum becomes too flat.

Conclusions

In this contribution, Monte Carlo studies on the acceleration of test particles in relativistic, subluminal shock environments are presented. Source candidates are Active Galactic Nuclei with boost




Figure 2: The diffuse energy spectrum of relativistic, subluminal sources, compared to the total diffuse spectrum.

factors of around $\Gamma \approx 10$ and Gamma Ray Bursts with highly-relativistic shocks, $100 < \Gamma < 1000$. The resulting particle spectra from those shocks have been investigated in the context of varying Γ and obliquity, i. e. the inclination angle between the shock normal and the magnetic field, ψ . For three angles, $\psi = 23^\circ, 33^\circ, 43^\circ$, shocks were simulated with nine different boost factors, $\Gamma = 10, 20, 30, 100, 300, 500, 700, 900, 1000$. While the result stayed constant for the different inclination angles, a linear behavior between the spectral index α_p and the boost factor Γ was found, leading to spectra of around $\alpha_p \sim 2.2$ for mildly-relativistic shocks ($\Gamma \sim 10$), but producing much harder spectra for highly-relativistic shocks. For boost factors around $\Gamma \sim 300 - 1000$, the spectra can become as flat as $\alpha_p \sim 1.5$.

Relativistic, subluminal shock occur in the jets of AGN ($\Gamma \sim 10$) and GRBs ($100 < \Gamma < 1000$). Since highly-relativistic shocks produce very flat spectra, the observed Cosmic Ray spectrum at the highest energies cannot be explained using a significant contribution from GRBs. A pure AGN spectrum on the other hand fits the spectrum well.

Acknowledgments

The project is co-funded by the European Social Fund and National Resources (EPEAEK II) PYTHAGORAS. Part of the work was also supported by the DFG grant LU1495/1-1.

References

- [1] W. I. Axford, E. Leer and G. Skadron, 1978, 15th ICRC Plovdiv, Bulgaria, Proc. 11, 132
- [2] M. C. Begelman and J. G. Kirk, 1990, ApJ 353, 66
- [3] A. R. Bell, 1978a, MNRAS 182, 147
- [4] A. R. Bell, 1978b, MNRAS 182, 443
- [5] R. D. Blandford and J. P. Ostriker, 1978, ApJL 221, L29
- [6] D. C. Ellison, F. C. Jones, S. P. Reynolds, 1990, ApJ 360, 702
- [7] D. C. Ellison, S. P. Reynolds, F. C. Jones, 1990 ApJ, 360, 702
- [8] G. Hasinger, T. Miyaji and M. Schmidt, 2005, APh 441, 417
- [9] D. P. Hudson 1965, MNRAS, 131, 23
- [10] F. de Hoffmann and E. Teller, 1950, Pr 80, 692
- [11] J. G. Kirk, P. Schneider, 1986 BAAS, 18, 908
- [12] G. F. Krymsky, 1977, Dokl.Acad.Nauk. SSSR 234, 1306
- [13] R. Lieu et al., 1994, ApJ, 421, 211
- [14] A. Meli, J. K. Becker and J. J. Quenby, 2007, to be submitted to A&A.
- [15] A. Meli and J. J. Quenby, 2003a, APh 19, 637
- [16] A. Meli and J. J. Quenby, 2003b, APh 19, 649
- [17] J. A. Peacock, 1981 MNRAS, 196, 135
- [18] R. J. Protheroe, A. Meli and A.-C. Donea, 2002, SpScRev 143, 276
- [19] J. J. Quenby, R. Lieu, 1989, Nature, 342, 654
- [20] A. Y. Gallant, A., Achterberg, J. K., Kirk, 1999, A&AS, 138, 549