



## Study of the 28 October 2003 solar flare by means of 2.223 MeV gamma-emission from it

E. V. TROITSKAIA<sup>1</sup>, L. I. MIROSHNICHENKO<sup>2</sup>

<sup>1</sup> D.V. Skobeltsyn Institute of Nuclear Physics (SINP), M.V. Lomonosov Moscow State University, 119992, Moscow, Russia

<sup>2</sup> Instituto de Geofísica, UNAM, C.U., Coyoacán, 04510, México, D.F., México  
troi@srd.sinp.msu.ru

**Abstract:** We have studied some characteristics of the 28 October 2003 powerful solar flare and surrounding medium by means of 2.223 MeV line time profile of gamma-emission. The modeling of time history and comparing it with the observational data reveals the considerable enhancement of the solar atmosphere density in the deep photospheric layers during the decay phase of gamma emission. Besides, the analysis let us detect the hardening of a neutron spectrum in the decay phase comparing to the rising one.

### Introduction

The neutron capture line of 2.223 MeV from solar flares has been analyzed directly or by using additional data to obtain the characteristics of particle acceleration in solar flares and properties of surrounding solar atmosphere. In particular, Ramaty and Murphy [13] used the fluences of 2.223 MeV line, for determining the energy spectra of solar energetic particles, Hua and Lingenfelter [7] studied their angular distributions and production of secondary neutrons, and Yoshimori et al. [17] investigated photospheric <sup>3</sup>He abundance.

The other authors [e.g. 10, 11] have developed an approach to determine the most probable profile of plasma density in the solar photosphere and adjoining levels during the period of a flare. By the 2.223 MeV line data on three large flares, 22 March 1991 [16], 6 November 1997 [11] and 16 December 1988 [12] the plausible model of the solar atmosphere density in the period of flare and some evidence of the effect of density enhancement was obtained. Gan [2, 3] was the first who applied the time profiles of calculated partial fluences of neutron capture line  $\varphi(2.223 \text{ MeV})$  and annihilation line  $\varphi(0.511 \text{ MeV})$  to deduce the spectral evolution of accelerated charged parti-

cles. Similar studies have been recently carried out for the RHESSI flare of 23 July 2002 [5].

In [7, 8] the energy spectra of solar flare neutrons were calculated in different suppositions on accelerated particle spectra. Founding on these neutron spectra, in the case of the flare 16 December 1988 the authors [12] deduced additionally the character of accelerated particles energy spectrum evolution.

In the present work we apply the developed approach to the powerful solar event of 28 October 2003.

### Method

The calculations of neutron propagation in the solar matter and 2.223 MeV line production are carried out using Monte-Carlo simulation, with due account for the models of vertical density profile of the solar plasma. For our SINP code we make allowance for: (1) neutron deceleration in elastic collisions with hydrogen nuclei, with due account for the energy and angular dependencies of cross-sections for  $np$ -scattering; (2) possible energetic neutron escape from the Sun; (3) gravitational neutron-Sun interaction; (4) thermal motion of decelerated neutrons; (5) neutron decay; (6) neutron captures by hydrogen <sup>1</sup>H, with the production of deuterium <sup>2</sup>H and gamma-quantum

of 2.223 MeV; (7) non-radiative neutron absorption on  $^3\text{He}$ ; (8) gamma-ray absorption in the solar atmosphere in dependence on solar flare central angle; (9) time profile of initial neutron production; (10) initial neutron spectra, and (11) altitude dependence of surrounding matter density. The relative abundance of  $^3\text{He}/\text{H}$  is taken about  $2 \times 10^{-5}$  [e.g., Hua and Lingenfelter, 1987; Yoshimori et al., 1999]. The time history of initial neutron production are assumed to be analogous (similar) to that of total fluence of  $^{12}\text{C}+^{16}\text{O}$  nuclear de-excitation lines in the range of 4.1-6.4 MeV. Calculations are made with SINP code for neutrons with energies of 1-100 MeV that are the most important ones for the 2.223 MeV line production. The primary neutrons are assumed to be emitted isotropically in the lower half-space (to-

wards the Sun) from the levels with densities less than  $5 \times 10^{15} \text{ cm}^{-3}$ . As a basic density model (BDM) ( $m=1$ ) we have used the standard astrophysical model HSRA (Harvard-Smithsonian Reference Atmosphere) for the lower chromosphere and quiet photosphere [6] together with a model of convection zone [14] consistent with the first one.

To determine possible deviations of the model, realizing in the observable flare from the BDM, we have also composed four additional models ( $m=2, 3, 4, 5$ ) representing smaller and larger densities at photospheric and adjoining levels as compared with the standard model ( $m=1$ ) of the quiet Sun (Table 1), see also e.g. [10]. In more detail, our method and calculation model (SINP code) are described elsewhere (see, e.g., [12].

Table 1. Characteristics of models.  $m$  is the number of model. Parameter  $\tau$  is the optical depth for a wavelength of 500 nm, the level  $\tau = 0.005$  corresponds to the top of the photosphere.

№, $m$	Character of density	
	Brief	Detailed
1	The main astrophysical model of the lower chromosphere and photosphere HSRA [12] together with model of convective zone of Spruit [13]	Grows gradually from $1.5 \cdot 10^{16} \text{ cm}^{-3}$ at the top of photosphere up to $2 \cdot 10^{17} \text{ cm}^{-3}$ at the 330 km level lower, where $\tau=1$ and sharply grows up to $\tau=10$ in the depth 60 km.
2	Enhanced into and under the photosphere	Enhanced to $8 \cdot 10^{17} \text{ cm}^{-3}$ at the depth $\sim 500$ km under the top of photosphere.
3	Enhanced into and under the photosphere	Grows up more slowly under the photosphere and mounts to $6 \cdot 10^{17} \text{ cm}^{-3}$ at the same depth.
4	Depressed from the low chromosphere down to the lower levels	Reduction of the density from the low chromosphere down to the deep layers. The density is $3 \cdot 10^{15} \text{ cm}^{-3}$ at the top of photosphere and $2 \cdot 10^{16} \text{ cm}^{-3}$ at the 330 km lower level.
5	Enhanced in the whole thickness of the photosphere.	The density is $2 \cdot 10^{17} \text{ cm}^{-3}$ through the complete thickness of the photosphere.

## Results

The flare of 28 October 2003 began at 9:41 UT, had its maximum at 11:10 and ended about 11:24 UT. It lasted about 15 min in the gamma-ray band. It appeared in the NOAA active region 10486. We apply our method to investigate the 28 October 2003 solar flare of X17.2/4B importance with coordinates S16E08 [15] and present the results for this powerful and long-duration

flare. The data on 2.223 MeV and summarized fluxes of 4.44 and 6.13 MeV gamma emission from INTEGRAL are used [9].

The calculations of time profiles of gamma fluxes were made in supposition of Bessel form (stochastic acceleration) of accelerated particles energy spectrum for three meanings of spectraparameter  $\alpha T$ : 0.005, 0.03 and 0,1 (Fig.1).

Table 2. The least square sums ( $\Sigma$ )

$\alpha T$	model	$\Sigma$
0.005	1	5.1031E+02
	2	5.3191E+02
	3	5.5098E+02
	4	6.3365E+02
	5	1.3819E+02
0.03	1	2.5450E+02
	2	2.3119E+02
	3	2.4751E+02
	4	2.9343E+02
	5	1.3002E+02
0.1	1	2.2200E+02
	2	1.9599E+02
	3	2.1556E+02
	4	2.5121E+02
	5	1.3227E+02

## Analysis

The table 1 shows that the best modeling time profile is in the case of  $\alpha T=0.03$  and  $m=5$ . This means the density enhancement in the whole thickness of photosphere. We can also conclude from the Fig. 1 that  $m=5$  begin to realize about 400 s. Another conclusion is that the better fitting in the rising phase is  $\alpha T=0.005$  and in the phases of maximum and decay the best fitting is  $\alpha T=0.1$

## Conclusion

In the present work we confirm the previously conclusions about the density enhancements in the deep photospheric or subphotospheric layers that were made for 3 flares. The hardening of particle spectra is also confirmed.

Since the character of 2.223 MeV gamma-emission time history depends mainly on the accelerated particles energy spectrum, density of surrounding atmosphere, content of  $^3\text{He}$  in the solar atmosphere and angular distribution of accelerated particles, it would be useful to have methods to reveal the portion of separate




Figure 1. Observational data of 2.223 MeV gamma-emission and modeling with spectral indexes 0.005, 0.03 and 0.1 for five models.

components. In present there appeared some appointments on the possible registration of the gamma-line at 20.58 MeV arising from radiative absorption neutrons on  $^3\text{He}$  [1].

## Acknowledgements

The authors are very indebted to Dr. V. Tatischeff for assignment the data on the solar flare of 28 October 2003.

This work has been supported partly by the Russian Foundation of Basic Research (RFBR, projects 02-02-39032, 03-02-96026, 05-02-39011), Federal Purpose Scientific and Technical Program (Section I, Project 4), and President's Grant of Russian Federation (project 1445.2003.2). The work of W. Gan was supported by NNSFC (China) via grants 10221001, 10333040 and by grant 2006CB806302 from the Ministry of Science and Technology of China. The work by L.I. Miroshnichenko was also supported partly by the CONACyT, Mexico (project 45822, PERPJ10332).

## References

- [1] Arkhangel'skaja I.V. et al. The observation of gamma-ray emission during January 20 2005 solar flare. In this volume.
- [2] Gan, W.Q. Spectral Evolution of Energetic Protons in Solar Flares. *Astrophys. J.* 496, 992-997, 1998
- [3] Gan, W.Q.: 2000, Energetic protons: spectral evolution and its significance, in R. Ramaty and N. Mandzhavidze (eds.), ASP Conf. Series, High Energy Solar Physics – Anticipating HESSI. 206, 439-444, 2000
- [4] Gan, W., Kuzhevskij, B.M., Miroshnichenko, L.I., and Troitskaia, E.V. Time profile of the 2.223 MeV line emission and some features of the 16 December 1988 solar event., in: Proc. ISCS 2003 Symposium, "Solar Variability as an Input to the Earth's Environment", Tatranska Lomnica, Slovakia, ESA SP-535, pp.655-657, 2003,
- [5] Gan, W.Q. On both the time histories of the 0.511 MeV line and 2.223 MeV line from the X4.8 flare of 23 July 2002 observed with RHESSI., *Solar Phys.* 219, 279-287, 2004.,
- [6] Gingerich, O., Noyes, R.W., Kalkofen, W., and Cuny, Y. The Harvard-Smithsonian reference atmosphere, *Solar Phys.* 18, 347-365, 1971.,
- [7] Hua, X.-M. and Lingenfelter, R.E. Solar flare neutron production and the angular dependence of the capture gamma-ray emission. *Sol. Phys.* 107, 351-383. 1987.
- [8] Hua, X.-M., Kozlovsky, B., Lingenfelter, R.E., Ramaty, R. and Stupp, A., Angular and energy-dependent neutron emission from solar flare magnetic loops, *Astrophys. J. Suppl.* **140**, 563-579. 2002
- [9] Kiener J., Gros M., Tatischeff V., and Weidenspointner A & A V. 445, P. 725-733, 2006
- [10] Kuzhevskij, B.M., Kuzhetsov, S.N., and Troitskaia, E.V. Development of the solar flare plasma density investigation method based on the 2.2 MeV gamma-line time profile analysis, *Adv. Space Res.* 22(7), 1141-1147, 1998.
- [11] Kuzhevskij, B.M., Miroshnichenko, L.I. and Troitskaia, E.V., Derivation of density profiles in the solar atmosphere by the 2.223 MeV line data for the 6 November 1997 flare, Proc. 27th Int. Cosmic Ray Conf., Invited, Rapporteur and Highlight Papers (Special Volume), 285-288, 2001.
- [12] Kuzhevskij, B.M., Miroshnichenko, L.I. and Troitskaia, E.V. Gamma-ray radiation with energy of 2.223 MeV and the density distribution in the solar atmosphere during flares, *Russian Astronomy Reports.* 49, 566-577 (in English), 2005
- [13] Ramaty, R. and Murphy, R.J. Nuclear processes and accelerated particles in solar flares *Space Sci. Rev.* 45(3/4), 213-268, : 1987
- [14] Spruit, H.C., A model of the solar convection zone. *Solar Phys.* 34, 277-290. 1974,
- [15] Veselovskiy I.S. et al. (75 authors) *Kosmicheskie Issledovaniya*, V.42, No 5, P.1-57, 2004.
- [16] Troitskaia E.V. and Kuzhevskij B.M. Absorption of 2.22 MeV solar flare gamma-rays and determining of the solar plasma density altitude profile, in Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA. August 17-25, 1999, 6, pp. 17-20, 1999.
- [17] Yoshimori, M., Shiozawa, A., and Suga, K. Photospheric  $^3\text{He}$  to H abundance ratio derived from gamma-ray line observations, Proc. 26th Int. Cosmic Ray Conf., 6, 5-8, 1999