



## Study of cosmic ray shower front and time structure with ARGO-YBJ

A.K. CALABRESE MELCARNE<sup>1,3</sup>, I.DE MITRI<sup>1,3</sup>, G. MARSELLA<sup>2,3</sup>, L. PERRONE<sup>2,3</sup>, G. PETRONELLI<sup>1,3</sup>, A.SURDO<sup>3</sup>, G.ZIZZI<sup>1,3</sup> ON BEHALF OF THE ARGO-YBJ COLLABORATION

<sup>1</sup>*Dipartimento di Fisica Università del Salento, Lecce, Italy*

<sup>2</sup>*Dipartimento di Ingegneria dell'Innovazione Università del Salento, Lecce, Italy*

<sup>3</sup>*INFN (Istituto Nazionale di Fisica Nucleare, Lecce, Italy)*

*giovanni.marsella@le.infn.it*

**Abstract:** ARGO-YBJ is a full coverage Extensive Air Shower array located by the high-altitude cosmic rays laboratory of Yangbajing (4300 m a.s.l., Tibet, China). The detector consists of a layer of Resistive Plate Counters covering an area of about 6500 m<sup>2</sup>; the time is measured on unit cell of 56 × 62 cm<sup>2</sup> with time resolution of about 1 ns. The high granularity of the detector allows a detailed characterization of cosmic ray showers. A set of well reconstructed data has been used to study the shower front structure. Simulated showers have been generated and the detector response has been taken into account in detail for this analysis. Several observables have been investigated in both real and simulated data and compared.

### Introduction

The ARGO-YBJ [1] experiment (Astrophysical Radiation with Ground-based Observatory at Yangbajing) has been designed to study cosmic rays and cosmic  $\gamma$ -radiation with energy greater than a few hundreds GeV, by detecting air showers at high altitude with wide-aperture and high duty cycle. The detector consists of a single layer of Resistive Plate Counters (RPCs), 280 × 125 cm<sup>2</sup> each, covering an area of 77.85 × 74.5 m<sup>2</sup> [2]. The percentage of active area in the central array is about 93%. To improve the reconstruction capability, the surrounding area has been partially ( $\sim 20\%$ ) instrumented with a guard ring of RPCs extending the detector layout up to 109.5 × 98.5 m<sup>2</sup>. A 0.5 cm thick lead converter will cover uniformly the RPC plane in order to improve the angular resolution. The detector is structured in 5.7 × 7.6 m<sup>2</sup> clusters, each consisting of twelve RPCs. A cluster is the basic detection and Data Acquisition logical unit. Pads of 55.6 × 61.8 cm<sup>2</sup> are the time elemental units ("pixels") for measuring the pattern of the shower front with time resolution of about 1 ns.

Ten pads cover one RPC and each pad contains 8 pick-up strips of 6.75 × 61.8 cm<sup>2</sup>, which provide a large particle counting dynamic range. The construction of the detector has finished. The central carpet (5800 m<sup>2</sup>, 130 clusters) is operating since July 2006 with a multiplicity trigger of  $N_{\text{pad}} > 20$ . The trigger rate is 4.1 kHz and the data flow is 7 MB/s. First results are reported in [3].

### Shower Reconstruction

The ARGO-YBJ digital readout allows detecting shower secondary particles down to very low densities. The time profile of the shower front can be reconstructed by the time of fired pads. Each pad time resolution is determined by the intrinsic performance of the RPC and its electronics and by signal propagation along the strips. Relative time offsets among different pads are corrected by a proper 'timing calibration' [4]. The overall time resolution is of about 1 ns. Particles within several tens of meters from the shower core are expected to form a curved front. Indeed, to reconstruct the primary particle arrival direction, space-time co-

ordinates (position and time of the fired pads in the event) can be fitted to a cone. A Maximum Likelihood based algorithm is used to perform a reliable reconstruction of the shower core position up to the edge and slightly beyond the active carpet [5]. The pad signals are shaped to 90 ns and sent to a multi-hit TDC so that particles hitting the same pad with a delay time larger than 90 ns are recorded up to a maximum delay of about  $1.3 \mu\text{s}$ .

## Time structure of extensive air shower front

The space-time structure of extensive air showers depends on primary mass, energy and arrival direction and on the interaction mechanisms with air nuclei. Measurements of shower parameters with several detection techniques would be required for a detailed knowledge of the shower front. A sketch illustrating the technique for studying an extensive air shower front with a surface detector is shown in Fig. 1. A flat array can measure the particles arrival times and their densities at ground.

Detailed studies of the shower front structure have been carried out by several groups and presented for example in [6, 7, 8] and related references.

The high space-time granularity of the ARGO-YBJ detector allows a fine sampling of the shower front close to the core. The time structure of the shower disk has been studied as a function of the distance to the shower axis up to 40 m in 1 m bins. The following observables have been investigated in the energy range between few TeV up to 20 TeV as a function of the distance to the shower axis:

- **the curvature** of the shower front as the mean of time residuals with respect to a planar fit ( $T_d$  in Fig. 1)
- **the thickness** of the shower front as the root mean square (RMS) of time residuals with respect to a conical fit ( $T_s$  in Fig. 1).

A set of well reconstructed data consisting of about  $10^6$  events has been selected for this




Figure 1: Sketch of shower front geometry and observables

analysis. The reconstructed cores are required to be within an area of  $20 \times 20 m^2$  centered on the carpet. A hit (pad) multiplicity greater than 200 and a quality cut on the  $\chi^2/\text{dof}$  of the fit has been applied in order to reject misreconstructed events. A Monte Carlo study has shown that the applied cuts select well reconstructed events with core inside the carpet (contained events).

Fig. 2 (left) shows the mean of time residuals with respect to a planar fit as a function of the distance to the shower axis for different pad multiplicities and for zenith angles less than  $15^\circ$ . It can be derived from simulation that hit multiplicities of 200, 400, 600 and 800 correspond to average primary energies of about 4, 7, 11 and 15 TeV (proton primaries, zenith angle less than  $15^\circ$ ). The deviation from a planar fit increases with distance (up to about 8 ns at 40 m) and depends only weakly on hit multiplicity in the considered energy range.

Fig. 2 (right) shows the RMS of time residuals with respect to a conical fit as a function of the distance to the shower axis for different pad multiplicities and for zenith angles less than  $15^\circ$ . The thickness of the shower front increases with distance (up to about 7 ns at 40 m) without any significant dependence on hit multiplicity in the considered energy range. Finally, no significant dependence on zenith angle has been observed both for curvature and thickness of the shower front up to  $45^\circ$ .




Figure 2: Mean of time residuals with respect to a planar fit (left) and RMS of time residuals with respect to a conical fit (right) as a function of distance to the shower axis for different pad multiplicities and for zenith angles less than  $15^\circ$ . Real data shown (statistical uncertainties only).

### Comparison of simulation with data

A Monte Carlo (MC) study has been performed using a sample of about  $6 \times 10^6$  CORSIKA showers [9] (protons) with zenith angle less than  $45^\circ$  and energy spectral index of  $-2.7$ , in the range between 100 GeV and 1000 TeV. The detector response has been simulated in detail with a GEANT3-based program [10]. A preliminary comparison between data and simulation is discussed.

Fig. 3 shows the distribution of pad multiplicity from simulation (solid line) in comparison with data (bullets), for zenith angle less than  $15^\circ$ . A fairly good agreement is found. Fig. 4 shows the time residuals with respect to a planar fit for data (blue bullets) and simulation (red boxes) (pad multiplicity between 200 and 400 and zenith angle less than  $15^\circ$ ). An agreement is found up to a distance of 15 m. At larger distances, the measured front curvature tends to be larger than predicted by simulation. Similarly, the measured shower thickness has been found to be systematically larger than the expectation and this effect is larger at small distances to the shower axis. As a final remark, it should be pointing out that this simulation does not include the contribution of heavier primary nuclei and some discrepancy between data and simulation remains for the




Figure 3: Distribution of pad multiplicity for data and simulation (MC). The histograms are normalized to the same area.

case of multiple hits. These facts may account for the differences observed in Fig. 3 at very high multiplicity.

To inspect the dependence of the studied observables on primary mass, a dedicated simulation has been performed for proton and photon showers. 5000 protons and 5000 photons showers have been generated in the energy range between 3 and 10 TeV with zenith angle between  $0^\circ$  and  $15^\circ$ . The RMS of time residuals with respect to a conical fit is shown in Fig. 5 as a function of the distance to the shower axis,




Figure 4: Time residuals with respect to a planar fit for data and simulation (MC), for pad multiplicity between 200 and 400 and zenith angle less than  $15^\circ$ .

for proton and photon primaries. The selected events are required to have a pad multiplicities larger than 200 and reconstructed core within an area of  $40 \times 40 m^2$  centered on the carpet. Though the measured difference is very small (at the level of the time resolution of the detector), this observable could be investigated in order to provide, on statistical basis, a clue to  $\gamma/hadrons$  separation and mass composition studies. Further studies and an extended simulation with heavier primaries are planned.

## Conclusions

The space-time high-granularity of the ARGO-YBJ detector provides a powerful tool for the study of time structure of cosmic rays shower front close to the core. Curvature and thickness of the shower front have been measured in the energy range from few TeV up to 20 TeV. The impact of these observables on mass composition studies has been discussed and the potentialities of the method investigated.

## References

- [1] M. Abbrescia et al. (ARGO-YBJ Coll.), *Astroparticle Physics with ARGO*, Pro-




Figure 5: RMS of time residuals with respect to a conical fit for proton and photon primaries (pad multiplicity between 200 and 400 and zenith angle less than  $15^\circ$ ).

- posal (available at <http://argo.na.infn.it>), 1996.
- [2] G. Aielli et al. (ARGO-YBJ Collaboration), *Nucl. Instr. Meth.*, A562 (2006) 92.
- [3] D. Martello on behalf of the ARGO-YBJ Collaboration, these Proceedings.
- [4] H.H. He, P. Bernardini, A.K. Calabrese Melcarne and S.Z. Chen, *Astropart. Phys.* (doi:10.1016/j.astropphys.2007.03.004), physics/0701291, P. Bernardini et al., on behalf of the ARGO-YBJ Collaboration 29th ICRC 5 (2005) 147 (Pune, India).
- [5] G. Di Sciascio on behalf of the ARGO-YBJ Collaboration 28th ICRC (2003) Tsukuba, Japan.
- [6] J. Linsley and L. Scarsi, *Phys. Rev.* 128 (1962) 2384.
- [7] G. Agnetta et al. *Astropart. Phys.* 6 (1997) 301.
- [8] M. Ambrosio et al. *Astropart. Physics* 11 (1999) 437.
- [9] D. Heck et al., “*CORSIKA: A Monte Carlo Code to Simulate Extensive Air Showers*” Report FZKA 6019, (1998).
- [10] *GEANT - Detector Description and Simulation Tool* CERN Program Library Long Writeup W5013 (1994).