**30th International Cosmic Ray Conference** 



## A Spectacular VHE Gamma-ray Outburst from PKS 2155-304 in 2006

W. BENBOW<sup>1</sup>, C. BOISSON<sup>2</sup>, L. COSTAMANTE<sup>1</sup>, O. DE JAGER<sup>3</sup>, D. EMMANOULOPOULOS<sup>4</sup>,

B. GIEBELS<sup>5</sup>, S. PITA<sup>6</sup>, M. PUNCH<sup>6</sup>, C. RAUBENHEIMER<sup>3</sup>, M. RAUE<sup>7</sup>, H. SOL<sup>2</sup>, S. WAGNER<sup>4</sup>

<sup>1</sup> Max-Planck-Institut für Kernphysik, Heidelberg, Germany

<sup>2</sup> LUTH, UMR 8102 du CNRS, Observatoire de Paris, Section de Meudon, France

<sup>3</sup> Unit for Space Physics, North-West University, Potchefstroom, South Africa

<sup>4</sup> Landessternwarte, Heidelberg, Germany

<sup>5</sup> LLR, IN2P3/CNRS, Ecole Polytechnique, Palaiseau, France

<sup>6</sup> Astrophysique et Cosmologie (APC), Paris, France

<sup>7</sup> Universität Hamburg, Institut fuer Experimentalphysik, Germany

Wystan.Benbow@mpi-hd.mpg.de

**Abstract:** A spectacular outburst of VHE (>100 GeV)  $\gamma$ -ray emission was observed from PKS 2155–304 in 2006 with the HESS experiment. The VHE flux reached levels of  $\sim$ 20 times the Crab Nebula flux, more that 2 orders of magnitude above typical values observed from the object. This extremely high flux, when coupled with the sensitivity of HESS, enables temporal studies of the emission with unprecedented resolution. Several fast flares, analogous to bursts, were resolved allowing for very strong constraints on the underlying physical processes. The temporal characteristics, spectral evolution and broadband characteristics of these 2006 outbursts will be presented. In addition, the challenges these data present to standard paradigms of VHE Blazars, particle acceleration processes, and radiation mechanisms will be discussed.

Results will be given in the post-conference version of the proceedings.

berg, Palaiseau, Paris, Saclay, and in Namibia in the construction and operation of the equipment.

## Acknowledgements

The support of the Namibian authorities and of the University of Namibia in facilitating the construction and operation of H.E.S.S. is gratefully acknowledged, as is the support by the German Ministry for Education and Research (BMBF), the Max Planck Society, the French Ministry for Research, the CNRS-IN2P3 and the Astroparticle Interdisciplinary Programme of the CNRS, the U.K. Particle Physics and Astronomy Research Council (PPARC), the IPNP of the Charles University, the Polish Ministry of Science and Higher Education, the South African Department of Science and Technology and National Research Foundation, and by the University of Namibia. We appreciate the excellent work of the technical support staff in Berlin, Durham, Hamburg, Heidel-